



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

excluded it only remains to apply the test for determining whether in the proposed equations (5) and (6), the auxiliary cubic is necessary. The general quartic

$$x^4 + ax^3 + bx^2 + cx + d = 0$$

is solvable by quadratics, without the intervention of a cubic, as shown by Thomas Simpson, in his Algebra, in three cases, viz., when

$$c = \frac{1}{2}af, \text{ or } a_1/d \text{ or } 2_1/df,$$

where, for brevity, $f = b - \frac{1}{4}a^2$. Considering equation (6),

$$\begin{aligned} a &= +3 & d &= -4 \\ b &= -21 & f &= -23\frac{1}{4} \\ c &= -64 \end{aligned}$$

whence $\frac{1}{2}af = -34\frac{7}{8}$; $a_1/d = 3\sqrt{-5}$; $2_1/df = 5\sqrt{31}$.

As none of these quantities equal c , the conclusion is that y , and similarly x , cannot be determined by *strictly quadratic methods*.

Also solved by J. E. SANDERS, and G. B. M. ZERR.

GEOMETRY.

191. Proposed by J. V. ADAMS, St. Louis, Mo.

Trisect any angle by means of the hypocycloid.

Solution by G. B. M. ZERR, A. M., Ph. D., The Temple College, Philadelphia, Pa.; JOHN J. QUINN, Warren High School, Warren, Pa.; and M. E. GRABER, Heidelberg University, Tiffin, O.

Let EAE' be an arc of the hypocycloid from cusp to cusp; R —radius of fixed circle; r —radius of generating circle; AQB the central generating circle; O its center; Q any point on this circle. Join QB , QO . With QO as a radius and O as a center describe the arc QD meeting the hypocycloid in D . Let GDF be the generating circle when D is the generating point, GHE its diameter. Draw DF ; construct angle BOK —to angle ACQ . Now arc GD —arc AQ , arc BQ —arc DF —arc FE .

\therefore arc GD —arc BF —arc AQ . Arc BF measures angle BOF , arc AQ measures angle ACQ . But arc $BF = r/R$ arc BEK .

\therefore Angle $BOF = r/R$ angle $BOK = r/R$ angle ACQ . If $R = 3r$, angle $BOF = \frac{1}{3}$ angle ACQ .

Therefore, by means of a suitable hypocycloid any angle may be divided in any given ratio. This property is applicable to the epicycloid also.

Also solved by the PROPOSER.

